


The impact of Toll-like receptors on the immune system functioning and on the immunopathogenesis of chronic hepatitis C: a modern view (literature review)

L. M. Syzova *^{A-E}, T. I. Koval ^{A-F}, V. I. Ilchenko ^{B,C,D}, I. M. Zviaholska ^{B,C,D},
O. M. Iziumska ^{B,C,D}

Ukrainian Medical Stomatological Academy, Poltava

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation; D – writing the article; E – critical revision of the article; F – final approval of the article

Key words:

Toll-like receptors, polymorphism, chronic hepatitis C.

Pathologia

2020; 17 (1), 114-121

*E-mail:

isizof@gmail.com

There exists a considerable body of literature on immunopathogenesis of chronic hepatitis C. Although results appear consistent with prior research in the area mentioned above, they appear inconsistent with the issues in the area of diagnosis, prognosis and treatment effectiveness. In this context the study addresses the research to receptors of the innate immune system – Toll-like receptors.

The aim of the research is to analyze the data of current professional literature regarding the role of individual Toll-like receptors in the immunopathogenesis of chronic hepatitis C.

Materials and methods. The method of reviewing and systematizing as well as the method of content analysis were used to overview the scientific literature as for the role of Toll-like receptors. For this purpose, we employ survey data collected from the world professional literature and analyzed the results of current researches.

Conclusions. The innate immune system plays a prominent role in the primary protection of the body against pathogens which recognition depends on the Toll-like receptors family whereas the genetic analysis is considered as a promising method of preventive and personalized medicine. The advantage of genetic markers regardless of age and other factors contain information about the susceptibility to multifactorial diseases which can be used to create a «genetic passport» of a person. Perceptions about the impact of the Asp299Gly polymorphism of the Toll-like receptor 4 gene and Gln11Leu of the Toll-like receptor 7 gene on the immunopathogenesis of chronic hepatitis C are ambiguous and this research provides a good starting point for discussion and further study which will allow optimizing the therapeutic and diagnostic tactics for this disease based on the complex evaluation of immunity which are defined by the determined polymorphisms.

Ключові слова:

Toll-подібні рецептори, поліморфізм, хронічний гепатит С.

Патология. 2020.

Т. 17, № 1(48).
С. 114-121

Сучасний погляд на роль Toll-подібних рецепторів у функціонуванні імунної системи та імунопатогенезі хронічного гепатиту С (огляд літератури)

Л. М. Сизова, Т. І. Коваль, В. І. Ільченко, І. М. Звягольська, О. М. Ізіумська

Незважаючи на великі досягнення у вивченні імунопатогенезу хронічного гепатиту С, залишається багато невирішених питань щодо діагностики, прогнозування перебігу та ефективності лікування. У контексті вивчення цієї інфекції викликають інтерес рецептори вродженої імунної системи – Toll-подібні рецептори.

Мета роботи – проаналізувати відомості сучасної фахової літератури щодо ролі окремих Toll-подібних рецепторів в імунопатогенезі хронічного гепатиту С.

Матеріали та методи. Методами оглядового, системного та контент-аналізу опрацювали доступні наукові джерела, що присвячені вивченню Toll-подібних рецепторів. Наведені дані світової фахової літератури та проаналізовані результати сучасних досліджень.

Висновки. Вроджена імунна система відіграє визначальну роль у первинному захисті організму від патогенів, розпізнавання яких залежить від родини Toll-подібних рецепторів, а генетичний аналіз – перспективний метод превентивної та персоналізованої медицини. Перевага генетичних маркерів полягає в тому, що вони незалежно від віку та інших факторів містять інформацію про схильність до мультифакторіальних хвороб, що можна використати під час створення «генетичного паспорта» людини. Уявлення щодо впливу поліморфізму Asp299Gly гена Toll-подібного рецептора 4 і Gln11Leu гена Toll-подібного рецептора 7 на імунопатогенез хронічного гепатиту С неоднозначні та потребують продовження вивчення. Це дасть змогу оптимізувати лікувально-діагностичну тактику при цьому захворюванні на основі комплексного оцінювання особливостей імунного реагування, що зумовлені наявністю названих поліморфізмів.

Современный взгляд на роль Toll-подобных рецепторов в функционировании иммунной системы и иммунопатогенезе хронического гепатита С (обзор литературы)

Л. М. Сизова, Т. И. Коваль, В. И. Ильченко, И. Н. Звягольская, Е. М. Изюмская

Несмотря на значительные достижения в изучении иммунопатогенеза хронического гепатита С, остается много нерешенных вопросов относительно диагностики, прогнозирования течения и эффективности лечения. В кон-

тексте изучения этой инфекции вызывают интерес рецепторы врожденной иммунной системы – Toll-подобные рецепторы.

Цель работы – провести анализ данных современной научной литературы о роли отдельных Toll-подобных рецепторов в иммунопатогенезе хронического гепатита С.

Материалы и методы. Методами обзорного, системного и контент-анализа обработаны доступные научные источники, посвященные изучению Toll-подобных рецепторов. Представлены данные мировой научной литературы и проанализированы результаты современных исследований.

Выводы. Врожденная иммунная система играет ведущую роль в первичной защите организма от патогенов, распознавание которых зависит от семьи Toll-подобных рецепторов, а генетический анализ – перспективный метод превентивной и персонализированной медицины. Преимущество генетических маркеров заключается в том, что они независимо от возраста и других факторов содержат информацию о склонности к мультифакториальным заболеваниям, что может быть использовано при создании «генетического паспорта» человека. Представление о влиянии полиморфизма Asp299Gly гена Toll-подобного рецептора 4 и Gln11Leu гена Toll-подобного рецептора 7 на иммунопатогенез хронического гепатита С неоднозначны и требуют дальнейшего изучения, что позволит оптимизировать лечебно-диагностическую тактику при этом заболевании на основе комплексной оценки особенностей иммунного реагирования, обусловленных наличием указанных полиморфизмов.

Ключевые слова:

Toll-подобные рецепторы, полиморфизм, хронический гепатит С.

Патология. 2020.
Т. 17, № 1(48).
С. 114-121

Despite the great advances in the study of immunopathogenesis of chronic hepatitis C a number of questions regarding diagnosis, prognosis and effectiveness of treatment of this infection remain to be addressed. We present an overview of literature that relates to work presented here. In recent years there has been renewed interest in mechanisms of the damaging effect of hepatotropic viruses on the liver and the role of the immune system in the progression of pathology. This issue provokes a number of gaps and shortcomings. A thorough understanding of the immunopathogenetic features of chronic hepatitis C and the role of individual immune system receptors in this process will significantly improve chronic hepatitis C therapy and advance an individual treatment approach, minimizing the financial costs and harm of side effects of medicines. Future advancements of gene polymorphism investigations as a factor in genetic susceptibility to infectious diseases and their peculiarities are expected to provide new opportunities for identifying at-risk groups and selecting personalized therapy for each patient. The study of chronic hepatitis C, the Toll-like receptors (TLRs) is a topic under intense research and pivotal importance.

Aim

The aim of the research is to analyze the data of modern professional literature regarding the role of individual TLRs in the immunopathogenesis of chronic hepatitis C.

Materials and methods

In the course of the expertiment methods of review, system and content analysis allowed us to explore and delve into the study of TLRs. We present an overview of the world professional literature that relates to study presented here.

A recent study on the role of TLRs in the functioning of the immune system.

TLRs were first detected in *Drosophila melanogaster* in 1985 by C. Nüsslein-Volhard (German) and E. Wieschaus (USA), the discovery of TLRs was an important event for the research in immunology and was highly estimated, thus Hoffmann (Luxembourg) and B. Beutler (USA) were awarded the Nobel Prize in Physiology or

Medicine in 2011 [1]. TLR4 was the first element to be discovered, then other TLRs in mammals and humans came [2]. Currently 13 TLRs are known, and 10 TLRs of which have been studied in humans [3–4]. Most TLRs are located on the cell surface – TLR1, TLR2, TLR5, TLR6, TLR10; an example of an intracellular arrangement is TLR3, TLR7, TLR8, TLR9, some TLRs can be expressed both intracellularly and extracellularly (TLR4, TLR11, TLR12, TLR13) [2–3,5].

Receptors of the TLRs family recognize pathogens and, upon activation, increase the local synthesis of proinflammatory cytokines, prostaglandins, chemokines that trigger the mechanism of implementation of the inflammatory response – a cascade of adapter and signaling molecules that leads to the induction of innate and adaptive immunity [5–8]. TLRs interact with adapter molecules and transmit an immunogenic signal to effectors, transcription factors, and target genes that together form a TLR-dependent signaling pathway. This pathway functions as a complex, consistent system of functionally interacting molecules. The most conservative role of TLR-activation of antimicrobial immunity in the skin, mucous membranes of the respiratory, gastrointestinal and urogenital tracts [8–11].

The analysis of the functions of different TLRs revealed that cells of the innate immune system activate different signaling pathways depending on the infectious pathogen. The genes which encode elements of the signaling pathway are coherently regulated in concert. Genetic variability not only of TLRs but also of molecules of TLR-dependent signaling pathways can play an important role in recognizing PAMPs (pathogen-associated molecular structures) and altering the immune response to infection [3,8,12]. It is reasonable to claim that clarifying the role of TLR in infectious pathology will let on conducting diagnostics timely and predicting the nature of the disease in advance. Moreover, this allows to study the pathogenetic aspects of its development, as well as to justify the choice of adequate therapy [12–13].

TLR participation in innate immunity is ensured by:

- initiating of secretion of proinflammatory cytokines which are required for physiological immunological response under various influences, among which various infections occupy the central place [6];

– regulating of neutrophil activity; a special role is played by TLR2 and TLR4, when the first one mentioned above protects cells from apoptosis, and the second one manifests itself respectfully as an important regulator of neutrophil survival [14];

– controlling of activation, differentiation and survival of B-lymphocytes, in which TLR2, TLR4 and TLR9 take an active part (this pathway of activation of B-lymphocytes is accompanied by increased calcium emission, phosphorylation of some kinases, enhanced endocytosis, immunoglobulin synthesis, lymphocytes) [15];

– providing support for the intestinal immunity associated with the expression of TLR by the epithelial cells of its mucosa [10];

– participation in the functioning of cells of the central nervous system, most of which express TLR (microglia, neurons, astrocytes, endothelial cells of the brain vessels) and the differential impact of TLR on the function of microglia [16].

Equally important is the involvement of TLR in adaptive immunity, which is also accomplished through a number of mechanisms:

– activation of CD4- and CD8-T lymphocytes [17];

– stimulation of the functions of different antigen recognition dendritic cells that express TLR2, TLR3, TLR4, TLR7, TLR9 [18];

– activation of macrophages, mast cells, in particular, with the participation of TLR9, which is especially pronounced when exposed to the genetic material of DNA viruses, bacteria, fungi [19];

– regulation of homeostasis of fibroblasts, myofibroblasts, fibroblast-like synoviocytes, endothelial and epithelial cells, in particular, with the participation of TLR2, TLR4, TLR6 [20];

– regulation of normal epithelial cells (TLR2, TLR3, TLR4, TLR5) as well as endothelial cells [21–22];

– potentiation of adaptive immunity with the inclusion of different mechanisms [20].

TLR transmembrane proteins have an extracellular domain (leucine-rich repeat, LRR) and an intracellular Toll IL-1 receptor domain (TIR). The LRR domain recognizes bacterial patterns and transmits the signal. TIR domain is a conserved peptide that interacts with proteins. It is also a part of several cytoplasmic proteins, including two Myeloid differentiation primary response gene (MyD88 and TIR domain containing adapter protein – TIRAP). These proteins transmit TLR signals that may be general and specific. The first signals are induced from all, the second – from one particular type of TLR.

Common signaling pathways are represented by adapter proteins (MyD88 and Toll-interacting protein – TOLLIP), protein kinases (IL-1R-associated kinase – IRAK) and TNF-receptor associated factor 6 (TRAF-6) adapter protein. The interaction between these structures leads to the activation of a large family of mitogen-activated protein kinases (MAPKs). The entire signaling pathway of TLR activation is complex and multicomponent, but the final stage of the protein kinase reaction cascade is the activation of transcriptional factors in the cytosol of the cell in a locked (inactive) state [23]. Several groups of transcription factors are known today. However, the nuclear factor (NF- κ B) has been widely studied [8]. The value

of protein kinases is in the following. On releasing from the NF- κ B blocker, it enters the nucleus of the cell, where it binds to the promoter regions of the inducible gene, which in turn leads to its activation and the initiation of the synthesis of molecules which are encoded by the formation of specific RNAs, and by the activation of inflammatory responses, including cytokine genes [5, 10].

Various macromolecules, including lipids, carbohydrates, proteins, and nucleic acids, act as microbial ligands for TLR. The most well-known TLR ligands of exogenous origin include the components of the cell wall of bacteria: peptidoglycans, lipopolysaccharides, flagellin, DNA of microorganisms, viral RNA and many others. It has now been proven that TLR ligands can also be endogenous molecules released by cell necrosis and massive tissue destruction, as well as by the breakdown of extracellular matrix molecules [8, 24]. TLR binding to ligands triggers an inflammatory signaling cascade through cytoplasmic TIR domains involving adapter proteins, namely MyD88, TIRAP, TIR-domain-containing adapter-inducing interferon- β (TRIF), which leads to the production of cytokines, antimicrobial peptides and it is realized by intracellular signaling in two possible ways [23]. Typically, TLR, with the exception of TLR3, use the MyD88-dependent pathway in which IRAK is sequentially activated, which includes 4 subunits – 2 active kinases (IRAK-1 and IRAK-4) and 2 non-catalytic subunits (IRAK-2 and IRAK-M). Then IRAK-4 phosphorylates IRAK-1. Phosphorylated IRAK-1 binds to TRAF-6 and simultaneously activates NF- κ B, mitogen-activated protein kinases (MAPKs). In the case of NF- κ B, the protein remains in the cytoplasm, binding to the NF- κ B (κ B) inhibitor. Phosphorylation of κ B the κ B kinase complex (IKK) leads to the destruction of the inhibitor, which makes it possible to move NF- κ B to the nucleus. Simultaneous activation of NF- κ B and MAPKs induces transcription of various inflammatory genes, including tumor necrosis factor (TNF), interleukins (IL) 1, 6, 8 and 12. In addition, TLR4 can trigger an immune response via the MyD88-independent pathway. It is carried out by an adapter protein capable of inducing interferon (IFN), which leads to phosphorylation of transcription factors: interferon-regulating factor-3 (IRF-3) and NF- κ B. These two transcription factors potentiate the action of each other, stimulate the production of type I interferons (IFN- α , IFN- β , IFN- γ) and activation of interferon-inducible genes [8, 23–24]. The TLR7, TLR8, TLR9 genes are also capable of initiating type I interferon synthesis through the MyD88-dependent signaling pathway upon activation of IRF-3 and IRF-7 [5, 24]. In addition, signals from TLR7, activating IRF-7, stimulate the production of antiviral cytokines, including type I and II interferon, which in turn would activate the processes of destruction of intracellular pathogens, while activation of NF- κ B induces a pro inflammatory effect through cytokine secretion, such as TNF- α , IL-6 and IL-12 [25–26]. General scheme of TLRs signaling pathways by J. Howell et al. (2013) is depicted in the Fig. 1 [26].

The wide range of TLRs ligands and their presence on most cells of the body contributes to the involvement of TLR in the pathogenesis of many diseases [5–6, 8, 12]. Defects in the TLRs gene system: impaired ligand recognition, TLR expression, signal transduction, production of

effector molecules, and their polymorphism can lead to the development of infectious, autoimmune and oncological diseases, allergopathology [2,5,8,27–30].

The point mutations at the DNA level that encode the structure of the receptor can disrupt its normal functioning and underlie the individual variability of the human genome [30]. The most common cause of differences in gene structure is so-called single nucleotide polymorphism (SNP), which occurs after about 290 base pairs, resulting in the formation of specific alleles of genes, which affects the development of protective reactions and susceptibility to particular diseases [5,8,11,28–30]. Nowadays, the conceptual basis of predictive genomic medicine is the very idea of genetic polymorphism [31].

The gene polymorphism implies that several variants can be copied from the same gene, structurally different from the copy of the same protein, some of them copied or not active at all, or may have the opposite function. When localized in exons, SNPs change the codons of the genetic code, which can lead to replacement of amino acids in peptides, sometimes their presence is associated with an increase or decrease in the concentration/activity of the gene product, thus genetic polymorphism is a source of individual differences, in particular in susceptibility and their course [28,32]. The presence of predisposition genes does not mean that a person can necessarily develop certain pathological conditions, gene polymorphism does not allow determining the time of the disease occurrence, but allows us to identify the characteristics of metabolism, drug metabolism and individual risk of susceptibility to a particular disease.

Thus, our findings indicate that in infectious pathology TLRs gene polymorphism affects the individual features of immunity and leads to immunological disorders caused by activating and inactivating genetic damage to the receptors of innate recognition or signaling molecules [5,8]. According to S. Mukherjee et al. (2019) a common feature of TLRs functional polymorphism is a decrease in the ability to recognize the corresponding ligands, which results in less pronounced cell activation after encountering pathogens [32]. It is reported that in infectious diseases, «mutations» of genes can lead to impaired recognition of infectious agents and imbalance of the functioning of the system of innate immunity, which will eventually be manifested by increased susceptibility to infections and susceptibility to the development of chronic inflammatory processes [5,8,13,28,30].

Thus, over the past several decades there has been a sustained research activity in TLRs genes area not only in terms of basic knowledge but also in the practicalities of predictive and personalized medicine [31]. Along there are some studies in the issue discussed but still the research about the role of TLRs gene polymorphism in chronic hepatitis C remains limited which makes the relevance of our study.

Impact of Asp299Gly polymorphism of TLR4 gene and Gln11Leu of TLR7 gene on chronic hepatitis C immunopathogenesis.

Nowadays, the rapid accumulation of knowledge about the genetic basis of pathological processes enriches the idea of immunopathogenesis of common human




Fig. 1. TLRs signaling pathways [26].

diseases, which, of course, belongs to chronic hepatitis C. In terms of studying both the immunopathogenesis of the disease the gene TLR4, which interacts with the protein envelope of viruses, structural and non-structural proteins of hepatitis C virus (HCV), and the TLR7 gene, a ligand of which is single-stranded are of particular interest [3,5,6,25,26].

The TLR4 gene is located in the 9th chromosome (9q32-33i), its exogenous ligands are lipopolysaccharides of gram-negative bacteria, lipoteichoic acid, taxol, flavolipin, F-protein of respiratory syncytial virus, type 1 fimbriae, and fimbriae, mycobacterium tuberculosis ligands, viral glycoproteins, and endogenous – fibronectin, low density lipoproteins, heat shock proteins, β -defensins, HMGB-1 hyaluronan [9,33]. The polymorphic marker Asp299Gly (rs4986790) of the TLR4 gene is a single nucleotide replacement of adenine (A) by guanine (G) at the +896 position of exon 3, which leads to amino acid replacement of aspartic acid (Asp) by glycine (Gly) chain 29. This missense mutation, altering the structure of the extracellular domain of TLR4, leads to the loss of negative charge of the site at position 299, which disrupts the process of recognition of bacterial lipopolysaccharide [4, 10].

According to the results of research studies, the frequency of point mutations of the TLR4 gene is very low (<1 %), with the exception of the genetic polymorphism Asp299Gly, the frequency of which is >5 % [8,28,34]; Asp299Gly is virtually undetectable in Asian populations [33]. The practical significance of this polymorphism is associated with inhibition of phosphorylation after lipopolysaccharide stimulation, which in turn leads to a decrease in the translocation of NF- κ B into the nucleus and affects the inhibition of the synthesis of proinflammatory cytokines IL-6 and TNF- α and production of IL-10 [33].

The TLR7 gene is located on Xp22.2. chromosomes and is a component of the body's antiviral defense system. It consists of a single exon and encodes a transmembrane protein of 1049 amino acids, it is localized in intracellular membrane compartments – endosomes, which in turn isolates it from a possible contact with endogenous nucleic

acids. The receptor is able to activate the transcription factors NF- κ B and IRF-7 [25]. The signal is transmitted through the adapter proteins MyD88, TRAF-6, IRAK-4 [35–36]. Exogenous ligands for TLR7 are small synthetic compounds, single-stranded RNA characteristic of the viral genome, nucleoside analogues (imidazoquinolines), loxoribine, bromyrimine, endogenous auto-RNA, ribonucleoproteins [23,25].

Gln11Leu polymorphism (rs179008) of the TLR7 gene is located in exon 3 and is a single nucleotide replacement of adenine (A) for thymine (T) which leads to the amino acid change of glutamine (Gln) for leucine (Leu) in the 11 codon of the protein, this allelic variant is one of the three SNPs that occur in populations of more than 5 % [26,36–37]. Actual data regarding to the incidence of Gln11Leu polymorphism in healthy populations differs both in general and in gender, which is related to the X chromosomal localization of this gene. For example, S. A. Taghavi et al. (2009) report that its overall prevalence is 14.67 %, among women – 10.3 %, among men – 16.24 % [35], and according to M. Bordignon et al. (2013) – 44.2 % for women and 18.5 % for men [29]. This polymorphism encodes functionally defective proteins and is able to reduce IFN- α production, thereby disrupting the adaptive immune response that occurs through the TLR7-dependent signaling pathway [29, 37].

Currently, a large number of genes have been correlated with susceptibility, features, and efficacy of chronic hepatitis C antiviral therapy. A great attention is paid to the role of TLR genes, which are important participants in CSF immunopathogenesis. There are reports in the world and national research studies that highlight the role of TLR4 and TLR7 genes and their functioning in the immunopathogenesis of chronic hepatitis C.

Scientific data suggest that HCV recognition by immune cells occurs through TLR4, and signals from this gene are capable of regulating HCV replication [38–39]. It is stated that in patients with chronic hepatitis C there is a significant increase in the expression of TLR4 by peripheral blood mononuclear cells regardless of HCV genotype or histological stage of the disease, which increases the synthesis of IFN- β , IL-6 from B cells of the immune system and is activated in the immune system. Increased expression of TLR4 by hepatocytes and epithelial cells of the biliary tract, a positive correlation of this phenomenon with liver damage in inflammation, activation of myofibroblasts and the development of liver fibrosis in chronic hepatitis C indicate the results of studies by S. Y. Mohamed et al. (2017) [40].

In some studies, the prevalence of the TLR4 gene Asp299Gly polymorphism among chronic hepatitis C patients is studied. Thus, in the works of O. D. S. Pires-Neto et al. (2015) and A. A. Al-Qahtani et al. (2014) it is indicated that the Asp299Gly polymorphism of the TLR4 gene was detected in 6.9–9.1 % of patients with chronic hepatitis C, whereas according to C. Guarner-Argente et al. (2010) it reaches 40.0 % [41–43]. In L. Sizova et al. (2016) it is claimed that the prevalence of Asp299Gly polymorphism among chronic hepatitis C patients is 15.2 % and exceeds population control data 4.5 times, which gives reason to consider its carrier as a risk factor for chronic hepatitis C infection with HCV [44].

A large number of existing studies show that carriers of the polymorphic genotype Asp299Gly of the TLR4 gene have a more severe course of chronic hepatitis C which means that the viral load registered is higher and susceptibility to bacterial infections is increased, which is associated with decreased production of IL-6 and IL-10 or with the development of encephalopathy and also the inflammation as a reaction which can block the translocation of bacterial antigens in cirrhosis [43,45–47]. However, in other papers, there is no significant influence of this polymorphism on the clinical and laboratory characteristics of chronic hepatitis C [48–50].

The scientific data on the effect of the TLR4 gene Asp299Gly polymorphism on the process of fibrosis in the liver and its rate in chronic hepatitis C are extremely controversial. Thus, a number of studies claim that it has a connection to a higher rate of progression of liver fibrosis and the risk of cirrhosis [43,51] as well, other studies, by contrast, claim that the ability of Asp299Gly can slow down the clinical progression of chronic hepatitis C and act as a protective factor for the development of cirrhosis and hepatocellular carcinoma [52,53], whereas A.A. Al-Qahtani et al. (2014), O. D. S. Pires-Neto et al. (2015), and G. Dubinskaya et al. (2016), independently of one another, concluded that there is no influence on this process [41–43,54].

The results of studies by M. Peris et al. (2015), G. M. Dubynska et al. (2016) and M. S. Iqbal et al. (2017) point to the association of the TLR4 gene Asp299Gly polymorphism with the low efficacy of interferon antiviral chronic hepatitis C therapy [46,55–56], but no similar effect was found in M. Emonts (2008) [49].

Scientific reports regarding the involvement of the TLR7 gene and its peculiarities in the pathogenesis of HCV infection are ambiguous. It is reported that through this gene the induction of innate immunity in HCV-infected is induced, the blocking of HCV RNA, the stimulation of production of antiviral cytokine IFN- α by dendritic cells [57–58], and in the presence of Gln11Leu polymorphism, TLR7 expression is reduced by hepatocytes with the lack of recognition of the virus, thereby limiting the production of IFN- α and IFN- λ , while the synthesis of IL-6 remains unchanged [39]. Another hypothesis concerns the ability of HCV to impair the expression and/or function of TLR7, whereas a higher level of expression of this gene by hepatocytes is observed in individuals with advanced liver fibrosis [59].

The prevalence of Gln11Leu polymorphism of TLR7 gene among chronic hepatitis C patients according to E. Ascar (2010) and E. Schott et al. (2007) generally exceeds 30.0 %, according to the research of S. A. Taghavi et al. (2009) is 15.2 %, and according to Y. S. Elsedawy et al. (2016) and T. I. Koval et al. (2018) it reaches 41.7 % and 30.9 % among women with chronic hepatitis C, and among men – 17.6 % and 14.4 %, respectively [35–37,60–61].

In the works of E. Ascar (2010) and L. M. Sizova (2016), no significant associations of this polymorphism were found with the clinical and laboratory characteristics of chronic hepatitis C [37,50].

Information on the effect of the TLR7 gene Gln11Leu polymorphism on the rate of progression of liver fibrosis is extremely controversial. Some authors point to the ab-

sence of associations between its presence and the progression of liver fibrosis in chronic hepatitis C [36–37,62], others to the protective role of polymorphism in this process in chronic hepatitis C [54], whereas in the study of F. Z. Fakhir (2018), this polymorphism is described as a profibrogenic factor [63].

E. Schott et al. (2008), M. El-Bendary et al. (2018) and S. I. Malov et al. (2018) report a correlation of the presence of the Gln11Leu polymorphism of the TLR7 gene with increased susceptibility to HCV and adverse response to interferon antiviral therapy in chronic hepatitis C patients [64–66], as confirmed by a study by S. A. Taghavi et al. (2009), but without the gender dependence of this fact [34], but it is denied in the works of Y. M. Mosaad et al. (2019) and S. S. Sleptsova et al. (2019) [62,67].

Thus, the analysis of scientific works of domestic and foreign authors shows that the susceptibility to HCV, as well as the formation of complications in humans are genetically determined and genes of the immune response and inflammation play an important role in this. In this regard, the creation of differential diagnostic and prognostic criteria for the course and consequences of chronic hepatitis C, developed on the basis of the study and analysis of the genetic material of each patient, is of particular importance.

Conclusions

1. The innate immune system plays a prominent role in the primary protection of the body against pathogens whose recognition depends on the TLRs family, and genetic analysis is a promising method of preventive and personalized medicine

2. The advantage of genetic markers is that, regardless of age and other factors, they carry information about the susceptibility to multifactorial diseases, which can be used to create a «genetic passport».

3. Perceptions about the impact of the Asp299Gly polymorphism of the TLR4 gene and Gln11Leu of the TLR7 gene on chronic hepatitis C immunopathogenesis are ambiguous and require further study, which will allow optimizing the therapeutic and diagnostic tactics for this disease based on a comprehensive evaluation of the features of immunoreactivity.

Prospects for further research. Future studies could fruitfully explore this issue further by studying the immunopathogenetic mechanisms of TLR4 impact as well as TLR7 genes impact on the course of chronic hepatitis C and other infectious pathologies.

Conflicts of interest: authors have no conflict of interest to declare.
Конфлікт інтересів: відсутній.

Надійшла до редакції / Received: 02.02.2020
Після доопрацювання / Revised: 23.02.2020
Прийнято до друку / Accepted: 26.02.2020

Information about authors:

Syzova L. M., MD, PhD, Associate Professor of the Department of Infectious Diseases with Epidemiology, Ukrainian Medical Stomatological Academy, Poltava.
ORCID ID: 0000-0002-8335-3295

Koval T. I., MD, PhD, DSc, Associate Professor, Head of the Department of Infectious Diseases with Epidemiology, Ukrainian Medical Stomatological Academy, Poltava.
ORCID ID: 0000-0003-2685-8665

Ilchenko V. I., MD, PhD, Associate Professor of the Department of Endocrinology with Childhood Infectious Diseases, Ukrainian Medical Stomatological Academy, Poltava.
ORCID ID: 0000-0002-1451-442X

Zviaholska I. M., MD, PhD, Associate Professor of the Department of Microbiology, Virology and Immunology, Ukrainian Medical Stomatological Academy, Poltava.
ORCID ID: 0000-0001-7531-3966

Iziumska O. M., MD, PhD, Associate Professor of the Department of Infectious Diseases with Epidemiology, Ukrainian Medical Stomatological Academy, Poltava.
ORCID ID: 0000-0002-9480-8988

Відомості про авторів:

Сизова Л. М., канд. мед. наук, доцент каф. інфекційних хвороб з епідеміологією, Українська медична стоматологічна академія, м. Полтава.

Коваль Т. І., д-р мед. наук, доцент, зав. каф. інфекційних хвороб з епідеміологією, Українська медична стоматологічна академія, м. Полтава.

Ільченко В. І., канд. мед. наук, доцент каф. ендокринології з дитячими інфекційними хворобами, Українська медична стоматологічна академія, м. Полтава.

Звягольська І. М., канд. мед. наук, доцент каф. мікробіології, вірусології та імунології, Українська медична стоматологічна академія, м. Полтава.

Ізюмська О. М., канд. мед. наук, доцент каф. інфекційних хвороб з епідеміологією, Українська медична стоматологічна академія, м. Полтава.

Сведения об авторах:

Сизова Л. М., канд. мед. наук, доцент каф. инфекционных болезней с эпидемиологией, Украинская медицинская стоматологическая академия, г. Полтава.

Коваль Т. И., д-р мед. наук, доцент, зав. каф. инфекционных болезней с эпидемиологией, Украинская медицинская стоматологическая академия, г. Полтава.

Ильченко В. И., канд. мед. наук, доцент каф. эндокринологии с детскими инфекционными болезнями, Украинская медицинская стоматологическая академия, г. Полтава.

Звягольская И. Н., канд. мед. наук, доцент каф. микробиологии, вирусологии та иммунологии, Украинская медицинская стоматологическая академия, г. Полтава.

Изюмская Е. М., канд. мед. наук, доцент каф. инфекционных болезней с эпидемиологией, Украинская медицинская стоматологическая академия, г. Полтава.

References

- [1] Volchenkov, R., Sprater, F., Vogelsang, P., & Appel, S. (2012). The 2011 Nobel Prize in physiology or medicine. *Scand J Immunol.*, 75(1), 1-4. <https://doi.org/10.1111/j.1365-3083.2011.02663.x>
- [2] Takeda, K., & Akira, S. (2015). Toll-Like receptors. *Current Protocols in Immunology*, 2015, 14.12.1-14.12.10. <https://doi.org/10.1002/0471142735.im1412s109>
- [3] Sasai, M., & Yamamoto, M. (2013). Pathogen Recognition Receptors: Ligands and Signaling Pathways by Toll-Like Receptors. *International Reviews of Immunology*, 32(2), 116-133. <https://doi.org/10.3109/08830185.2013.774391>
- [4] Dowling, J. K., & Mansell, A. (2016). Toll-like receptors: the swiss army knife of immunity and vaccine development. *Clinical & Translational Immunology*, 5, Article UNSP e85. <https://doi.org/10.1038/cti.2016.22>
- [5] Vijay, K. (2018). Toll-like receptors in immunity and inflammatory diseases: Past, present, and future. *International Immunopharmacology*, 59, 391-412. <https://doi.org/10.1016/j.intimp.2018.03.002>
- [6] Kawai, T., & Akira, S. (2011). Toll-like Receptors and Their Crosstalk with Other Innate Receptors in Infection and Immunity. *Immunity*, 34(5), 637-650. <https://doi.org/10.1016/j.immuni.2011.05.006>
- [7] Lester, S. N., & Li, K. (2014). Toll-Like Receptors in Antiviral Innate Immunity. *Journal of Molecular Biology*, 426(6), 1246-1264. <https://doi.org/10.1016/j.jmb.2013.11.024>
- [8] Medvedev, A. E. (2013). Toll-Like Receptor Polymorphisms, Inflammatory and Infectious Diseases, Allergies, and Cancer. *Journal of Interferon and Cytokine Research*, 33(9), 467-484. <https://doi.org/10.1089/jir.2012.0140>

- [9] Izmailova, O. (2018). Analysis of association between the polymorphic variants of the TLR2 (RS5743708) and TLR4 (RS4986790, RS4986791) genes with the health status of apparently healthy individual. *Problemy ekologiy ta medytsyny*, 22(1-2), 17-22. <https://doi.org/10.31718/mep.2018.22.1-2.03>
- [10] Kordjazay, N., Haj-Mirzaian, A., Rohani, M. M., Gelfand, E. W., Rezaei, N., & Abdolghaffari, A. H. (2018). Role of toll-like receptors in inflammatory bowel disease. *Pharmacological Research*, 129, 204-215. <https://doi.org/10.1016/j.phrs.2017.11.017>
- [11] Lebedeva, O. P., & Qirko, R. (2018). Ekspresiya toll-podobnykh retseptorov v zhenskoy reproduktivnom trakte i ee gormonal'naya regulyatsiya (obzor) [Expression of TOLL-like receptors in the female reproductive tract and its hormone regulation (review)]. *Research Results in Biomedicine*, 4(3), 3-17. [in Russian]. <https://doi.org/10.18413/2313-8955-2018-4-3-0-1>
- [12] Netea, M. G., Wijmenga, C., & O'Neill, L. A. J. (2012). Genetic variation in Toll-like receptors and disease susceptibility. *Nature Immunology*, 13(6), 535-542. <https://doi.org/10.1038/ni.2284>
- [13] Tartey, S., & Takeuchi, O. (2017). Pathogen recognition and Toll-like receptor targeted therapeutics in innate immune cells. *International Reviews of Immunology*, 36(2), 57-73. <https://doi.org/10.1080/08830185.2016.1261318>
- [14] Hacine-Gherbi, H., Denys, A., Carpentier, M., Heysen, A., Duflot, P., Lanos, P., & Allain, F. (2017). Use of Toll-like receptor assays for the detection of bacterial contaminations in icodextrin batches released for peritoneal dialysis. *Toxicology Reports*, 4, 566-573. <https://doi.org/10.1016/j.toxrep.2017.10.004>
- [15] Pettengill, M. A., van Haren, S. D., Li, N., Dowling, D. J., Bergelson, I., Jans, J., Ferwerda, G., & Levy, O. (2016). Distinct TLR-mediated cytokine production and immunoglobulin secretion in human newborn naive B cells. *Innate Immunity*, 22(6), 433-443. <https://doi.org/10.1177/1753425916651985>
- [16] Gesuete, R., Kohama, S. G., & Stenzel-Poore, M. P. (2014). Toll-Like Receptors and Ischemic Brain Injury. *Journal of Neuropathology and Experimental Neurology*, 73(5), 378-386. <https://doi.org/10.1097/nen.0000000000000068>
- [17] Saxena, M., & Bhardwaj, N. (2018). Re-Emergence of Dendritic Cell Vaccines for Cancer Treatment. *Trends in Cancer*, 4(2), 119-137. <https://doi.org/10.1016/j.trecan.2017.12.007>
- [18] Jimenez-Dalmaroni, M. J., Gerswhin, M. E., & Adamopoulos, I. E. (2016). The critical role of toll-like receptors – From microbial recognition to autoimmunity: A comprehensive review. *Autoimmunity Reviews*, 15(1), 1-8. <https://doi.org/10.1016/j.autrev.2015.08.009>
- [19] Wang, S., Charbonnier, L. M., Rivas, M. N., Georgiev, P., Li, N., Gerber, G., Bry, L., & Chatila, T. A. (2015). MyD88 Adaptor-Dependent Microbial Sensing by Regulatory T Cells Promotes Mucosal Tolerance and Enforces Commensalism. *Immunity*, 43(2), 289-303. <https://doi.org/10.1016/j.immuni.2015.06.014>
- [20] Khakpour, S., Wilhelmsen, K., & Hellman, J. (2015). Vascular endothelial cell Toll-like receptor pathways in sepsis. *Innate Immunity*, 21(8), 827-846. <https://doi.org/10.1177/1753425915606525>
- [21] Dmytriyev, D. V., & Nazarchuk, O. A. (2015). Suchasni uiavlennia pro rol Toll-podobnykh retseptoriv v imunofiziologichnomu homeostazi ta pry patolohii [Modern views on the role of Toll-like receptors in immune and physiological homeostasis and pathology]. *Aktualni Problemy Suchasnoi Medytsyny Visnyk Ukrainiskoi Medychnoi Stomatologichnoi Akademii*, 15(3, part 2), 313-319. [in Ukrainian].
- [22] Stewart, E. A., Wei, R. X., Branch, M. J., Sidney, L. E., & Amoaku, W. M. (2015). Expression of Toll-like receptors in human retinal and choroidal vascular endothelial cells. *Experimental Eye Research*, 138, 114-123. <https://doi.org/10.1016/j.exer.2015.06.012>
- [23] Petrov, R. V., Haitov, R. M., & Chereshev, V. A. (2017). Fiziologiya immunoj sistemy: kletochnye i molekulyarno-biologicheskie mekhanizmy [Physiology of the Immune System: Cellular and Molecular-Biological Mechanisms]. In M. A. Ostrovsky (Ed.), *100 Years of Physiology in Russia on the 100th Anniversary of the Pavlov Physiological Society* (pp. 96-119). Russian Foundation for Basic Research. [in Russian]. <https://doi.org/10.22204/2410-4639-2017-094-025-96-119>
- [24] Kawasaki, T., & Kawai, T. (2014). Toll-like receptor signaling pathways. *Frontiers in Immunology*, 5, 461. <https://doi.org/10.3389/fimmu.2014.00461>
- [25] Funk, E., Kottlitz, S., Gilliam, B., & Talwani, R. (2014). Ticking the TLR7 to cure viral hepatitis. *Journal of Translational Medicine*, 12, Article 129. <https://doi.org/10.1186/1479-5876-12-129>
- [26] Howell, J., Angus, P., Gow, P., & Visvanathan, K. (2013). Toll-like receptors in hepatitis C infection: Implications for pathogenesis and treatment. *Journal of Gastroenterology and Hepatology*, 28(5), 766-776. <https://doi.org/10.1111/jgh.12170>
- [27] Tizaoui, K., Kaabachi, W., Hamzaoui, K., & Hamzaoui, A. (2015). Association of Single Nucleotide Polymorphisms in Toll-like Receptor Genes With Asthma Risk: A Systematic Review and Meta-analysis. *Allergy Asthma & Immunology Research*, 7(2), 130-140. <https://doi.org/10.4168/air.2015.7.2.130>
- [28] Skevaki, C., Pararas, M., Kostelidou, K., Tsakris, A., & Routsias, J. G. (2015). Single nucleotide polymorphisms of Toll-like receptors and susceptibility to infectious diseases. *Clinical and Experimental Immunology*, 180(2), 165-177. <https://doi.org/10.1111/cei.12578>
- [29] Bordignon, M., Bargagli, E., Agostini, C., Cinetto, F., Baldo, V., Alaibac, M., & Rottoli, P. (2013). TLR7 Gln11Leu single nucleotide polymorphism in patients with sarcoidosis. *Sarcoidosis Vasculitis and Diffuse Lung Diseases*, 30(2), 157-161.
- [30] Vesnina, L. E., Izmailova, O. V., Shlykova O. A., & Kidashev, I. (2015). Features of NF- κ B-mediated signaltransduction and development of systemic inflammation in patients with diseases of internal organs are determined by microbial factor and individual reactivity of the body (review of own research findings). *Problemy ekologiy ta medytsyny*, 19(3-4), 30-37.
- [31] David, S. P. (2017). *Genomic and Precision Medicine: Primary Care: Third Edition*. London: Academic Press.
- [32] Mukherjee, S., Huda, S., & Babu, S. P. S. (2019). Toll-like receptor polymorphism in host immune response to infectious diseases: A review. *Scandinavian Journal of Immunology*, 90(1), Article UNSP e12771. <https://doi.org/10.1111/sji.12771>
- [33] Ziakas, P. D., Prodromou, M. L., El Khoury, J., Zintzaras, E., & Mylonakis, E. (2013). The Role of TLR4 896 A > G and 1196 C > T in Susceptibility to Infections: A Review and Meta-Analysis of Genetic Association Studies. *Plos One*, 8(11), Article UNSP e81047. <https://doi.org/10.1371/journal.pone.0081047>
- [34] Rudenko, K. A., Tuguz, A. R., Anokhina, E. N., & Muzhenya D. V. (2014). Polimorfizmy genov Toll-podobnykh retseptorov, assotsirovannyeh s nasledstvennoj otogoshennostyu i vozrastom manifestatsii bronhialnoj astmy [Gene polymorphisms of Toll-like receptors associated with hereditary abnormalities and age of asthma onset]. *Immunologiya*, 35(3), 160-163. [in Russian].
- [35] Taghavi, S. A., Damangir, H., Kamali Sarvestani, E., & Eshraghian, A. (2009). Relation between C. 32A> T polymorphism in TLR7 and response to treatment in chronic HCV-infection. *Armaghane danesh*, 14(2), 105-116.
- [36] Schott, E., Witt, H., Neumann, K., Taube, S., Oh, D. Y., Schreier, E., Vierich, S., Puhl, G., Bergk, A., Halang, J., Weich, V., Wiedenmann, B., & Berg, T. (2007). A Toll-like receptor 7 single nucleotide polymorphism protects from advanced inflammation and fibrosis in male patients with chronic HCV-infection. *Journal of Hepatology*, 47(2), 203-211. <https://doi.org/10.1016/j.jhep.2007.03.021>
- [37] Askar, E., Ramadani, G., & Mihm, S. (2010). Toll-Like Receptor 7 rs179008/Gln11Leu Gene Variants in Chronic Hepatitis C Virus Infection. *Journal of Medical Virology*, 82(11), 1859-1868. <https://doi.org/10.1002/jmv.21893>
- [38] Mei, S., Burchell, J., Skinner, N., Millen, R., Matthews, G., Hellard, M., Dore, G. J., Desmond, P. V., Sundararajan, V., Thompson, A. J., Visvanathan, K., & Sasadeusz, J. (2016). Toll-like Receptor Expression and Signaling in Peripheral Blood Mononuclear Cells Correlate With Clinical Outcomes in Acute Hepatitis C Virus Infection. *Journal of Infectious Diseases*, 214(5), 739-747. <https://doi.org/10.1093/infdis/jiw235>
- [39] Lee, C. M., Hu, T. H., Lu, S. N., Wang, J. H., Hung, C. H., Chen, C. H., & Yen, Y. H. (2016). Peripheral blood toll-like receptor 4 correlates with rapid virological response to pegylated-interferon and ribavirin therapy in hepatitis C genotype 1 patients. *Bmc Gastroenterology*, 16, Article 73. <https://doi.org/10.1186/s12876-016-0492-6>
- [40] Mohamed, S. Y., Mostafa, E. F., Hanafy, A. S., Atia, H., Metwally, A., & Marei, A. M. (2017). The relationship between expression of Toll-like receptor 4 in chronic hepatitis C patients and different stages of liver fibrosis. *Gastroenterology and Hepatology from Bed to Bench*, 10(4), 278-283. <https://doi.org/10.22037/gfbb.v0i0.1079>
- [41] Pires-Neto, O. D., de Sa, K. S. G., Santana, B. B., Gomes, S. T. M., Amoras, E. D. G., Conde, S. R. D., Demachki, S., Azevedo, V. N., Martins-Feitosa, R. N., Ishak, M. D. G., Ishak, R., & Vallinoto, A. C. R. (2015). Lack of Association between Polymorphisms of the TLR4 Gene and Infection with the Hepatitis B and C Viruses. *Mediators of Inflammation*, Article 150673. <https://doi.org/10.1155/2015/150673>
- [42] Al-Qahtani, A. A., Al-Anazi, M. R., Al-Zoghaibi, F., Abdo, A. A., Sana, F. M., Khan, M. Q., Albenmoussa, A., Al-Ashgar, H. I., & Al-Ahdal, A. N. (2014). The Association of Toll-Like Receptor 4 Polymorphism with Hepatitis C Virus Infection in Saudi Arabian Patients. *Biomed Research International*, Article 357062. <https://doi.org/10.1155/2014/357062>
- [43] Guamer-Argente, C., Sanchez, E., Vidal, S., Roman, E., Concepcion, M., Poca, M., Sanchez, D., Juarez, C., Soriano, G., & Guamer, G. (2010). Toll-like receptor 4 D299G polymorphism and the incidence of infections in cirrhotic patients. *Alimentary Pharmacology & Therapeutics*, 31(11), 1192-1199. <https://doi.org/10.1111/j.1365-2036.2010.04291.x>
- [44] Sizova, L., Koval, T., Kaidashev, I., Ilchenko, V., & Dubinskaya, G. (2016). Rol geneticheskogo polimorfizma Toll-like retseptorov 4 i 7 v razvitii khronicheskogo gepatita s i gendernye osobennosti ikh raspredeleniya [The role of genetic polymorphisms Toll-like receptor 4 and 7 in the chronic hepatitis c and gender features of their distribution]. *Georgian Medical News*, (250), 51-56. [in Russian].

- [45] Nieto, J. C., Sanchez, E., Roman, E., Vidal, S., Oliva, L., Guarner-Argente, C., Poca, M., Torras, X., Juarez, C., Guarner, C., & Soriano, G. (2014). Cytokine production in patients with cirrhosis and TLR4 polymorphisms. *World Journal of Gastroenterology*, 20(46), 17516-17524. <https://doi.org/10.3748/wjg.v20.i46.17516>
- [46] Peric, M., Bosnjak, Z., Sarkanj, B., Barbic, J., Antolovic-Pozgajn, A., Ruzman, N., Roksandic-Krizan, I., & Vukovic, D. (2015). Polymorphisms of Toll-like receptors 2 and 4 in chronically infected hepatitis C patients from north-east Croatia. *Archives of Virology*, 160(1), 297-304. <https://doi.org/10.1007/s00705-014-2283-0>
- [47] Pinero, P., Juanola, O., Caparros, E., Zapater, P., Gimenez, P., Gonzalez-Navajas, J. M., Such, J., & Frances, R. (2017). Toll-like receptor polymorphisms compromise the inflammatory response against bacterial antigen translocation in cirrhosis. *Scientific Reports*, 7, Article 46425. <https://doi.org/10.1038/srep46425>
- [48] Alvarado-Tapias, E., Guarner-Argente, C., Oblitas, E., Sanchez, E., Vidal, S., Roman, E., Concepcion, M., Poca, M., Gely, C., Pavel, O., Nieto, J. C., Juarez, C., Guarner, C., & Soriano, G. (2018). Toll-like receptor 4 polymorphisms and bacterial infections in patients with cirrhosis and ascites. *World Journal of Hepatology*, 10(1), 124-133. <https://doi.org/10.4254/wjh.v10.i1.124>
- [49] Emonts, M. (2008). *Polymorphisms in immune response genes in infectious diseases and autoimmune diseases*. (Doctoral Dissertation). Erasmus University Rotterdam.
- [50] Sizova, L. M. (2016). Osoblyvosti kliniko-laboratorykh kharakterystyk khronichnoho hepatytu S u khvorykh z polimorfizmom Asp299Gly henu TLR4 ta Gln11Leu henu TLR7 [Features of clinical and laboratory characteristics of chronic hepatitis C in patient with polymorphism Asp-299Gly of TLR4 gene and Gln11Leu of TLR7 gene]. *Aktualni problemy suchasnoi medytyny*, 16(4), 182-187. [in Ukrainian].
- [51] Zhu, Q., Zou, L., Jagavelu, K., Simonetto, D. A., Huebert, R. C., Jiang, Z. D., ... Shah, V. H. (2012). Intestinal decontamination inhibits TLR4 dependent fibronectin-mediated cross-talk between stellate cells and endothelial cells in liver fibrosis in mice. *Journal of Hepatology*, 56(4), 893-899. <https://doi.org/10.1016/j.jhep.2011.11.013>
- [52] Guo, J., & Friedman, S. L. (2010). Toll-like receptor 4 signaling in liver injury and hepatic fibrogenesis. *Fibrogenesis and Tissue Repair*, 3(1) <https://doi.org/10.1186/1755-1536-3-21>
- [53] Guo, J. S., Loke, J., Zheng, F., Hong, F., Yea, S., Fukata, M., Tarocchi, M., Abar, O. T., Huang, H. J., Sninsky, J. J., & Friedman, S. L. (2009). Functional Linkage of Cirrhosis-Predictive Single Nucleotide Polymorphisms of Toll-like Receptor 4 to Hepatic Stellate Cell Responses. *Hepatology*, 49(3), 960-968. <https://doi.org/10.1002/hep.22697>
- [54] Dubinskaya, G., Sizova, L., Koval, T., Kovalyova, E., & Kaydashv, I. (2016). Clinical and genetic predictors and prognostic model of rapidly progressive hepatic fibrosis in chronic hepatitis C. *Georgian Medical News*, (256-257), 37-45.
- [55] Dubinskaya, G. M., Koval, T. I., Sizova, L. M., Izyumskaya, E. M., Kotelevskaya, T. M., Sviridenko, N. P., Voloshina, L. G., & Voitenko, L. L. (2016). Analiz effektivnosti protivovirusnoi terapii khronicheskogo gepatita C u patsientov Poltavskoi oblasti [Analysis of antiviral therapy effectiveness for chronic hepatitis C patients in Poltava region]. *Klinicheskaya infektologiya i parazitologiya*, 5(4), 440-448. [in Russian].
- [56] Iqbal, M. S., Ashfaq, U. A., Khaliq, S., Masoud, M. S., Qasim, M., Haque, A., Ghani, M. U., & Jahan, S. (2017). Toll-like receptor 4 polymorphism as pretreatment predictor of response to HCV genotype 3a interferon-based treatment. *Future Virology*, 12(12), 738-745. <https://doi.org/10.2217/fvl-2017-0076>
- [57] Zhang, Y. L., Guo, Y. J., Li, B., & Sun, S. H. (2009). Hepatitis C virus single-stranded RNA induces innate immunity via Toll-like receptor 7. *Journal of Hepatology*, 51(1), 29-38. <https://doi.org/10.1016/j.jhep.2009.03.012>
- [58] Dominguez-Molina, B., Machmach, K., Perales, C., Tarancon-Diez, L., Gallego, I., Sheldon, J. L., Leal, M., Domingo, E., & Ruiz-Mateos, E. (2018). Toll-Like Receptor 7 (TLR-7) and TLR-9 Agonists Improve Hepatitis C Virus Replication and Infectivity Inhibition by Plasmacytoid Dendritic Cells. *Journal of Virology*, 92(23), Article e01219-18. <https://doi.org/10.1128/jvi.01219-18>
- [59] Dehghan-Manshadi, M., Hadinedoushan, H., Amirbaigy, M. K., Zare, F., Eslami, G., Mirghanizade-Bafghi, S. A., & Akhondi-Meybodi, M. (2015). Relative Expression of Toll-Like Receptors 2 and 7 mRNA in Peripheral Blood of Patients With Hepatitis C. *Hepatitis Monthly*, 15(11), Article e30427. <https://doi.org/10.5812/hepatmon.30427>
- [60] Elsedawy, Y. S., Khattab, M. A., & El Hady, S. A. (2016). Single Nucleotide Polymorphisms of Toll-Like Receptor 7 in Hepatitis C Virus Infection and Hepatocellular Carcinoma Patients. *The Egyptian Journal of Medical Microbiology*, 25(4), 73-80. <https://doi.org/10.12816/0037023>
- [61] Koval, T. I., Syzova, L. M., Dubynska, H. M., Pryimenko, N. O., Rudenko, S. S., Marchenko, O. H., & Shlykova, O. (2018). TLR4 and TLR7 genetic polymorphism in patients with HIV/HCV-coinfection: prevalence and gender-related features. *Wiadomosci Lekarskie (Warsaw, Poland : 1960)*, 71(8), 1566-1570.
- [62] Mosaad, Y. M., Metwally, S. S., Farag, R. E., Lofly, Z. F., & Abdel-Twab, H. E. (2019). Association between Toll-Like Receptor 3 (TLR3) rs3775290, TLR7 rs179008, TLR9 rs352140 and Chronic HCV. *Immunological Investigations*, 48(3), 321-332. <https://doi.org/10.1080/08820139.2018.1527851>
- [63] Fakhir, F. Z., Lkhider, M., Badre, W., Alaoui, R., Meurs, E. F., Pineau, P., ... Benjelloun, S. (2018). Genetic variations in toll-like receptors 7 and 8 modulate natural hepatitis C outcomes and liver disease progression. *Liver International*, 38(3), 432-442. <https://doi.org/10.1111/liv.13533>
- [64] Schott, E., Witt, H., Neumann, K., Bergk, A., Halangck, J., Weich, V., ... Berg, T. (2008). Association of TLR7 single nucleotide polymorphisms with chronic HCV-infection and response to interferon- α -based therapy. *Journal of Viral Hepatitis*, 15(1), 71-78. <https://doi.org/10.1111/j.1365-2893.2007.00898.x>
- [65] El-Bendary, M. E., Neamatallah, M., Elalfy, H., Besheer, T., Elkholi, A., El-Diasty, M. E., Elsareef, M., Zahran, M., El-Aarag, B. E., Gomaa, A., Elhamady, D., El-Setouhy, M. E., Hegazy, A., & Esmat, G. (2018). The association of single nucleotide polymorphisms of Toll-like receptor 3, Toll-like receptor 7 and Toll-like receptor 8 genes with the susceptibility to HCV infection. *British Journal of Biomedical Science*, 75(4), 175-181. <https://doi.org/10.1080/09674845.2018.1492186>
- [66] Malov, S. I., Baatarkhuu, O., Garkov, O. B., Ariunaa, S., Baigalmaa, J., Davaakhuu, B., Munk-Orshikh, D., Ouynbileg, J., Stepanenko, L. A., Savilov, E. D., Zherbanova, L. V., Symbelova, T. A., & Malov, I. V. (2018). The Role of Human Genetic Factors in the Natural Selection of Hepatitis C Virus' Dominant Genotype in Ethnically Close Populations of Buryats and Khalkha-Mongols. *Sovremennye Tehnologii V Meditsine*, 10(3), 21-30. <https://doi.org/10.17691/stm2018.10.3.3>
- [67] Sleptsova, S. S., Bilyukina, I. F., Malov, S. I., Semenov, S. I., Stepanenko, L. A., Ogarkov, O. B., Savilov, Y. D., & Malov, I. V. (2019). Polymorphism of genes localized on the X-chromosome as a marker of predisposition to infectious process chronicity at hepatitis C in the ethnic group of male Yakuts. *Yakut Medical Journal*(2), 64-67. <https://doi.org/10.25789/ymj.2019.66.19>