

Official journal of the Polish Medical Association



Memory of dr Władysław Biegański

Since 1928

VOLUME LXXIII, ISSUE 12 PART 1, DECEMBER 2020

Wiadomości Lekarskie is abstracted and indexed in: PUBMED/MEDLINE, SCOPUS, EMBASE, INDEX COPERNICUS, POLISH MINISTRY OF SCIENCE AND HIGHER EDUCATION, POLISH MEDICAL BIBLIOGRAPHY

Copyright: © ALUNA Publishing House.

Articles published on-line and available in open access are published under Creative Common Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Wiadomości Lekarskie monthly journal

You can order the subscription for the journal from Wydawnictwo Aluna by:

prenumerata@wydawnictwo-aluna.pl Wydawnictwo Aluna Z.M. Przesmyckiego 29 05-510 Konstancin-Jeziorna Poland

Place a written order first.

If you need, ask for an invoice.
Payment should be done to the following account of the Publisher:
account number for Polish customers (PLN):
82 1940 1076 3010 7407 0000 0000
Credit Agricole Bank Polska S. A., SWIFT: AGRIPLPR

account number for foreign customers (EURO): 57 2490 0005 0000 4600 7604 3035 Alior Bank S. A.: SWIFT: ALBPPLPW

Subscription of twelve consecutive issues (1-12): Customers in Poland: 360 PLN/year Customers from other countries: 320 EURO/year



Editor in-Chief:

Prof. Władysław Pierzchała

Deputy Editor in-Chief:

Prof. Aleksander Sieroń

Statistical Editor:

Dr Lesia Rudenko

Managing Editor:

Agnieszka Rosa - amarosa@wp.pl

International Editorial Office:

Lesia Rudenko (editor) – I.rudenko@wydawnictwo-aluna.pl Nina Radchenko (editor's assistant)

- n.radchenko@wydawnictwo-aluna.pl

Polish Medical Association (Polskie Towarzystwo Lekarskie):

Prof. Waldemar Kostewicz – President PTL

Prof. Jerzy Woy-Wojciechowski – Honorary President PTL Prof. Tadeusz Petelenz

Prof. Tadeusz Petelenz

International Editorial Board - in-Chief:

Marek Rudnicki

Chicago, USA

International Editorial Board – Members:

| Kris Bankiewicz | San Francisco, USA | George Krol | New York, USA |
|---------------------|-------------------------|---------------------|--------------------|
| Christopher Bara | Hannover, Germany | Krzysztof Łabuzek | Katowice, Poland |
| Krzysztof Bielecki | Warsaw, Poland | Henryk Majchrzak | Katowice, Poland |
| Zana Bumbuliene | Vilnius, Lithuania | Ewa Małecka-Tendera | Katowice, Poland |
| Ryszarda Chazan | Warsaw, Poland | Stella Nowicki | Memphis, USA |
| Stanislav Czudek | Ostrava, Czech Republic | Alfred Patyk | Gottingen, Germany |
| Jacek Dubiel | Cracow, Poland | Palmira Petrova | Yakutsk, Russia |
| Zbigniew Gasior | Katowice, Poland | Krystyna Pierzchała | Katowice, Poland |
| Andrzej Gładysz | Wroclaw, Poland | Tadeusz Płusa | Warsaw, Poland |
| Nataliya Gutorova | Kharkiv, Ukraine | Waldemar Priebe | Houston, USA |
| Marek Hartleb | Katowice, Poland | Maria Siemionow | Chicago, USA |
| Roman Jaeschke | Hamilton, Canada | Vladyslav Smiianov | Sumy, Ukraine |
| Andrzej Jakubowiak | Chicago, USA | Tomasz Szczepański | Katowice, Poland |
| Oleksandr Katrushov | Poltava, Ukraine | Andrzej Witek | Katowice, Poland |
| Peter Konturek | Saalfeld, Germany | Zbigniew Wszolek | Jacksonville, USA |
| Jerzy Korewicki | Warsaw, Poland | Vyacheslav Zhdan | Poltava, Ukraine |
| Jan Kotarski | Lublin, Poland | Jan Zejda | Katowice, Poland |
| | | | |

Distribution and Subscriptions:

Bartosz Guterman prenumerata@wydawnictwo-aluna.pl

Graphic design / production:

Grzegorz Sztank www.red-studio.eu

Publisher:

ALUNA Publishing House ul. Przesmyckiego 29, 05-510 Konstancin – Jeziorna www.wydawnictwo-aluna.pl www.wiadomoscilekarskie.pl www.wiadlek.pl

FOR AUTHORS

- The monthly "Wiadomości Lekarskie" Journal is the official journal of the Polish Medical Association. Original studies, review papers as well as case reports are published.
- 2. The publication of the manuscript in "Wiadomości Lekarskie" is paid. The cost of publishing the manuscript is PLN 1,000 plus 23% VAT (for foreign authors: since July 2021 250 Euro). If the first author of the manuscript is a member of the Editorial Board or a team of journal reviewers, we do not charge a fee for printing the manuscript, and if she or he is the next co-author the fee is PLN 500 plus 23% VAT. The publisher issues invoices. The fee should be paid after receiving positive reviews, and before publishing the manuscript. Membership of the Polish Medical Association with documented paid membership fees for the last 3 years is also the exempt from publication fee.
- Only papers in English are accepted for publication. The editors can help in finding the right person for translation or proofreading.
- 4. Papers should be sent to the editor via the editorial panel (Editorial System), available on the journal's website at https://www.wiadlek.pl. In order to submit an article, free registration in the system is necessary. After registration, the author should follow the instructions on the computer screen.
- 5. All editorial work is under control and using the editorial panel. This applies in particular to sending manuscripts, correspondence between the editor and author and the review process. In special cases, the editor may agree to contact outside the panel, especially in case of technical problems.
- 6. Acceptable formats for individual elements of the article are as follows:
 - A) Content of the article doc, docx, rtf, odt.
 - B) Tables doc, docx, rtf, odt
 - C) Figures JPG, GIF, TIF, PNG with a resolution of at least 300 dpi
 - D) Captions for figures and tables.
 - These elements are sent to the editor separately using the editorial panel. References and article metadata such as titles, keywords, abstracts etc. are supplemented by the author manually in the editorial panel in appropriate places.
- 7. The volume of original papers including figures and references must not exceed 21,600 characters (12 pages of typescript), and review papers up to 28,800 characters (16 pages).
- The original manuscript should have the following structure: Introduction, Aims, Material
 and methods, Results, Discussion and Conclusions which cannot be a summary of the manuscript.
- 9. When using abbreviations, it is necessary to provide the full wording at the first time they are used
- 10. In experimental manuscripts in which studies on humans or animals have been carried out, as well as in clinical studies, information about obtaining the consent of the Ethics Committee should be included.
- 11. The Editorial Board follow the principles contained in the Helsinki Declaration as well as in the Interdisciplinary Principles and Guidelines for the Use of Animals in Research, Testing and Education, published by the New York Academy of Sciences Ad Hoc Committee on Animal Research. All papers relating to animals or humans must comply with ethical principles set out by the Ethics Committee.
- 12. The abstract should contain 150-250 words. Abstracts of original, both clinical and experimental, papers should have the following structure: Aims, Material and methods, Results, Conclusions. Do not use abbreviations in the title or the abstract. The abstract is pasted or rewritten by the authors into the appropriate field in the application form in the editorial panel.
- 13. Keywords (3-5) should be given according to MeSH (Medical Subject Headings Index Medicus catalogs http://www.nim.nih.gov.mesh/MBrower.html). Keywords cannot be a repetition of the title of the manuscript.
- 14. Illustrative material may be black and white or color photographs, dearly contrasting or drawings carefully made on a white background. With the exception of selected issues, the Journal is printed in shades of gray (black and white illustrations).
- 15. The content of the figures, if present (e.g. on the charts), should also be in English
- 16. Links to all tables and figures (round brackets) as well as references (square brackets) the author must place in the text of the article.

- 17. Only references to which the author refers in the text should be included in the list of references ordered by citation. There should be no more than 30 items in original papers and no more than 40 items in review papers. Each item should contain: last names of all authors, first letters of first names, the title of the manuscript, the abbreviation of the journal title (according to Index Medicus), year, number, start and end page. For book items, please provide: author's (authors') last name, first letter of the first name, chapter title, book title, publisher, place and year of publication. It is allowed to cite websites with the URL and date of use of the article, and if possible the last names of the authors. Each literature item should have a reference in the text of the manuscript placed in square brackets, e.g. [1], [3–6]. Items should be organized as presented in Annex 1 to these Regulations.
- 18. When submitting the article to the editor, the authors encloses a statement that the work was not published or submitted for publication in another journal and that they take full responsibility for its content, and the information that may indicate a conflict of interest, such as:
 - financial dependencies (employment, paid expertise, consulting, ownership of shares, fees),
 - 2. personal dependencies,
 - 3. academic and other competition that may affect the substantive side of the work,
 - sponsorship of all or part of the research at the stage of design, collection, analysis and interpretation of data, or report writing.
- 19. The authors in the editorial panel define their contribution to the formation of scientific work according to the following key:
 - A Work concept and design
 - B Data collection and analysis
 - C Responsibility for statistical analysis
 - D Writing the article
 - E Critical review
 - F Final approval of the article.
- 20. In the editorial panel along with the affiliation, the author also gives her or his ORCID number.
- 21. The Journal is reviewed in double, blind review mode. The submitted papers are evaluated by two independent reviewers and then qualified for publishing by the Editor-in-Chief. Reviews are anonymous. The authors receive critical reviews with a request to correct the manuscript or with a decision not to qualify it for publishing. The procedure for reviewing articles is in line with the recommendations of the Ministry of Science and Higher Education contained in the paper "Good practices in review procedures in science" (Warsaw 2011). Detailed rules for dealing with improper publishing practices are in line with COPE guidelines. The publishing review rules are in the Review Rules section.
- 22. Each manuscript is subject to verification in the anti-plagiarism system.
- 23. Manuscripts are sent for the author's approval. The author's corrections should be sent within the time limit indicated in the system. No response within the given deadline is tantamount to the author's acceptance of the submitted material. In special cases, it is possible to set dates individually.
- Acceptance of the manuscript for publishing means the transfer of copyright to the Aluna Publishing House (Aluna Anna Łuczyńska, NIP 5251624918).
- 25. Articles published on-line and available in open access are published under Creative Common Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.
- 26. The authors receive a free PDF of the issue in which their mansucript is enclosed, and on request a printed copy. The printed copy is sent to the address indicated by the authors as the correspondence address.
- 27. Manuscripts not concordant with the above instructions will be returned to be corrected.
- 28. The editors do not return papers which have not been commissioned.
- 29. The editors take no responsibility for the contents of the advertisements.



CONTENTS

| CONTENTS | |
|---|------|
| ORIGINAL ARTICLES Aidyn G. Salmanov, Taras P. Bondar, Yaroslav V. Shkorbotun, Evelina A. Chumak, Volodymyr O. Shkorbotun, Olena V. Dementieva, Vadim V. Pererva PREVALENCE OF NASAL CARRIAGE OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS AMONG HEALTHCARE WORKERS IN THE DEPARTMENTS OF OTORINOLARYNGOLOGY AND DENTISTRY IN KYIV, UKRAINE | 2563 |
| Anatoly A. Avramenko INFLUENCE OF PECULARITIES OF ANATOMICAL STRUCTURE (SHAPE) OF DUODENAL BULB ON THE MECHANISM OF FORMATION OF "KISSING" ULCERS OF PATIENTS WITH CHRONIC NON-ATROPHIC GASTRITIS | 2568 |
| Marlena Kwiatkowska, Inga Chomicka, Jolanta Malyszko COMPARISON OF MOLECULAR AND SEROLOGICAL TESTS FOR SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2 (SARS-COV-2) IN POST-EXPOSURE EMPLOYEES OF THE NEPHROLOGY DEPARTMENT | 2572 |
| Viktoria V. Alekseeva, Alina S. Nechiporenko, Andrii V. Lupyr; Nadiia O. Yurevych, Vitaliy V. Gargin A METHOD OF COMPLEX EVALUATION OF MORPHOLOGICAL STRUCTURE OF OSTIOMEATAL COMPLEX COMPONENTS, LOWER WALL OF MAXILLARY AND FRONTAL SINUSES | 2676 |
| Lyudmyla V. Pakharenko, Yuri P. Vdovichenko, Nataliia Ya. Kurtash, Iryna O. Basiuha, Inna V. Kravchuk, Volodymyr D. Vorobii, Olena M. Kusa ESTRADIOL BLOOD LEVEL AND ESR1 GENE POLYMORPHISM IN WOMEN WITH PREMENSTRUAL SYNDROME | 2581 |
| Khrystyna R. Boriak, Lyudmyla E. Vesnina RELATIONSHIP OF THE EATING BEHAVIOR FEATURES WITH ANTHROPOMETRIC INDICATORS AND ENERGY VALUE OF THE DIET IN YOUNG PEOPLE WITH NORMAL WEIGHT AND OVERWEIGHT | 2586 |
| Tetiana I. Nimtsovych, Anatolii M. Kravchenko, Olha Yu. Mishcheniuk, Tamara Ya. Chursina, Kyrylo O. Mikhaliev, Viktor P. Polovyi VISIT-TO-VISIT BLOOD PRESSURE VARIABILITY AND TARGET ORGAN DAMAGE IN RURAL DWELLERS WITH UNCOMPLICATED ARTERIAL HYPERTENSION | 2591 |
| Piotr Szolc, Łukasz Niewiara, Marta Kawulak, Kornelia Szkodoń, Elzbieta Ostrowska-Kaim, Marcin Nosal, Marcin Krzanowski, Krzysztof Żmudka, Jacek Legutko, Bartłomiej Guzik NEUTROPHIL-LYMPHOCYTE RATIO AND PLATELET-LYMPHOCYTE RATIO AS PREDICTORS OF CORONARY MICROCIRCULATORY DISEASE OCCURRENCE AND OUTCOME IN PATIENTS WITH CHRONIC CORONARY SYNDROME AND NO SIGNIFICANT CORONARY ARTERY STENOSIS | 2598 |
| Olena L. Zolotukhina, Iuliia G. Romanova, Tetyana O. Pyndus, Georgy O. Romanov, Iryna M. Tkachenko EVALUATION OF EFFECTIVENESS OF TREATMENT-AND-PROPHYLACTIC COMPLEX IN TOBACCO-ADDICTED PATIENTS WITH CHRONIC GENERALIZED PERIODONTITIS ON THE BACKGROUND OF CHRONIC HYPERACID GASTRITIS | 2607 |
| Oleksij P. Kostyrenko, Nataliia I. Vynnyk, Mykhailo M. Koptev, Petro A. Hasiuk, Maksym I. Skrypnyk, Alevtyna M. Bilous, Serhii A. Proskurnya DENTAL CROWN BIOMINERALIZATION DURING ITS HISTOGENESIS | 2612 |
| Ołha M. Arustamian, Volodymyr S. Tkachyshyn, Vitalii E. Kondratiuk, Nataliia Yu. Tkachyshyna, Oleksandr Yu. Aleksiichuk, Irina V. Dumka INFLUENCE OF ADVERSE PRODUCTION FACTORS ON CEREBRAL HEMODYNAMIC IN LOCOMOTIVE DRIVERS | 2617 |
| Nikolay A. Scherbina, Liliya V Potapova, Shcherbyna Irina N. Scherbina, Oksana P. Lipko , Olga V. Mertsalova, Anastasiia A. Chekhunova MODERN METHODS OF COMPLEX CORRECTION OF PSYCHOSOMATIC DISORDERS IN PATIENTS WITH EXTERNAL GENITAL ENDOMETRIOSIS | 2623 |
| Monika Białkowska, Tomasz Stołtny, Jarosław Pasek, Michał Mielnik, Karol Szyluk, Kamil Baczyński, Robert Hawranek, Aleksandra Koczy-Baron, Sławomir Kasperczyk, Grzegorz Cieślar, Bogdan Koczy THE INFLUENCE OF HIP ARTHROPLASTY ON HEALTH RELATED QUALITY OF LIFE IN MALE POPULATION WITH OSTEOARTHRITIS HIP DISEASE | 2627 |
| Iryna V. Redka, Oksana Ya. Mikhalchuk, Zhama V. Sotnikova-Meleshkina ELECTROENCEPHALOGRAPHIC MONITORING OF THE BRAIN FUNCTIONAL CONDITION IN MODERN UKRANIAN SCHOOLCHILDREN | 2634 |
| Oleksii O. Holubenko, Anatolii F. Levytskyi, Oleksandr V. Karabenyuk RESULTS OF TREATMENT OF IDIOPATHIC CONGENITAL CLUBFOOT IN CHILDREN: A 3-YEAR FOLLOW UP STUDY | 2640 |

| Tamara M. Pasiieshvili, Natalia M. Zhelezniakova, Lyudmila M. Pasiyeshvili, Olga M. Kovalyova TOTAL ANTIOXIDANT ACTIVITY AND MANGANESE SUPEROXIDE DISMUTASE IN COMORBIDITY OF GASTROESOPHAGEAL REFLUX DISEASE AND AUTOIMMUNE THYROIDITIS IN STUDENT POPULATION | 2644 |
|---|------|
| Svitlana P. Yarova, Sergii O. Turchenenko, Iryna M. Tkachenko, Yuriy Yu. Yarov ANALYSIS OF THE DYNAMICS OF FUNCTIONAL OCCLUSION INDICATORS IN THE CONSTRUCTION OF OCCLUSION SURFACES OF ORTHOPEDIC STRUCTURES | 2651 |
| Svitlana P. Palamar, Halyna O. Vaskivska, Liudmyla L. Nezhyva, Liudmyla A. Nazarenko, Serhii S. Terekhov FORMATION OF EDUCATIONAL AND COGNITIVE COMPETENCY OF JUNIOR SPECIALISTS IN THE PROCESS OF SELF-EDUCATIONAL ACTIVITY: EXPERIENCIAL ASPECT | 2657 |
| Nataliia O. Riabushko, Roman M. Riabushko, Mykola M. Riabushko, Olena B. Riabushko ASSESSMENT OF QUALITY OF LIFE OF PATIENTS AFTER SURGERY MADE BYTHE IN-HOUSE DEVICES AND METHODS | 2664 |
| Yevhenii S. Khilinich, Vadym Yu. Davydenko, Ivan I. Starchenko, Mykhailo Ya. Nidzelskyi, Hanna M. Davydenko, Viktor V. Kuznetsov THE EFFECT OF MONOMER OF REMOVABLE DENTURE BASE RESIN ON THE STRUCTURAL ORGANIZATION OF THE GLANDULAR ZONE OF ALBINO RAT HARD PALATE IN THE EXPERIMENT | 2667 |
| REVIEW ARTICLES Oleksandr A. Halushko, Maryna A. Trishchynska, Tetiana M. Povietkina, Marianna V. Boliuk DIABETES MELLITUS IN COVID-19 PATIENTS: VERDICT OR NOT? | 2672 |
| Monika Adamczyk- Sowa, Barbara Gębka- Kępińska, Michał Kępiński MULTIPLE SCLEROSIS — RISK FACTORS | 2677 |
| Lyubov V. Smaglyuk, Maryna V. Trofymenko, Alevtyna M. Bilous MODERN ASPECTS IN DIAGNOSIS OF TONGUE`S PARAFUNCTIONS (LITERATURE REWIEV) | 2683 |
| CASE STUDY Liudmyla A. Vygivska, Olena A. Yakovenko, Evgen V. Blagoveschenskiy, Yevheniia M. Babadzhanian, Lesia A. Rudenko, Kyrylo V. Yakovenko CURRENT STATE OF THE PROBLEM OFABNORMAL INVASION OF THE PLACENTA IN OBSTETRICS | 2688 |

ORIGINAL ARTICLE



ANALYSIS OF THE DYNAMICS OF FUNCTIONAL OCCLUSION INDICATORS IN THE CONSTRUCTION OF OCCLUSION SURFACES OF ORTHOPEDIC STRUCTURES

10.36740/WLek202012117

Svitlana P. Yarova¹, Sergii O. Turchenenko¹, Iryna M. Tkachenko², Yuriy Yu. Yarov¹

¹DONETSK NATIONAL MEDICAL UNIVERSITY, KRAMATORSK, UKRAIN ²UKRAINIAN MEDICAL STOMATOLOGICAL ACADEMI, POLTAVA, UKRAINE

ABSTRACT

The aim: Is to evaluate the effectiveness of the proposed method of constructing occlusal surfaces of orthopedic structures by monitoring the dynamics of functional indicators of occlusion and the functional condition of the masticatory muscles.

Materials and methods: The object of the study was made up by 64 patients with included defects of the dentition of the 3rd class according to Kennedy. Patients were divided into two equal groups — the main group (patients were treated by using the suggested method of construction of occlusal surfaces of orthopedic structures) and control (dentition defects were replaced according to the traditional protocol).

Results: The results of evaluating the effectiveness of the proposed method showed that the index of symmetry of the anterior bundles of the temporal muscles dropped by 18.5%, the index of symmetry of the masticatory muscles reduced by 22%, the torque of the mandible reduced by 32%, the index of stabilizing occlusal — by 27%, relative efforts of balance disturbance for more than 10 % regarding the left and right reduced by a factor of 4.4, the rate of opening time was 1.5 times higher than before, the rate of time of opening of the jaws — 1.8 times.

Conclusions: The use of the suggested method of orthopedic treatment with temporary structures leads to probable changes in the studied functional parameters in comparison with the control group (p < 0.05).

KEY WORDS: occlusion relationship, masticatory muscle, orthopedic constructions

Wiad Lek. 2020;73(12 p. 1):2651-2656

INTRODUCTION

Despite the ever-increasing aesthetic demand of patients, the effective restoration of the function of the dental-maxillary system (DMS) [1-5] remains an important requirement for orthopedic structures. Therefore, special attention should be paid to occlusal disorders that occur in the result of orthopedic treatment, as well as partial loss of teeth, which is accompanied by changes in almost all the parts of the dental apparatus and can provoke a number of complications. The most common and dangerous of them is development of musculoskeletal disorders. Numerous clinical studies have shown that lesions of the temporomandibular joint (TMJ) are revealed in 70-80% of the population and rank third after caries and periodontal disease [6-9].

It is a well-known fact that the basis for the formation of physiological occlusion is equal distribution of functional loads in cases of sufficient number of correctly placed occlusal contacts and adequate placement of TMJ components [10-12]. Modern methods of functional assessment of the condition of the masticatory muscles and occlusal relationships allow reaching the most adequate balance of the thyroid gland objectively in real time [13-16]. In connection with the above-mentioned facts, we consider it appropriate

to develop the method of construction of occlusal surfaces of orthopedic structures which is based on monitoring and subsequent correction of the studied functional parameters of occlusion and masticatory muscle activity.

THE AIM

The aim is to evaluate the effectiveness of the proposed method of constructing occlusal surfaces of orthopedic structures by monitoring the dynamics of functional indicators of occlusion of T-Scan III Novus and the functional condition of the masticatory muscles BioEMG III with subsequent correction of the dynamic component of these parameters.

MATERIALS AND METHODS

The object of the study was made up by 64 patients with included defects of the dentition of the 3rd class according to Kennedy, not complicated by dental deformities, aged from 18 to 60 years. Criteria for inclusion of patients in the study were the following: the presence of informed consent to the study, violation of the integrity of the dentition in the area of premolars and molars, indications for permanent

prosthetics. Criteria for exclusion of patients from the study were: the presence of removable prostheses, the presence of orthopedic structures based on dental implants, periodontal disease with signs of masticatory muscle hyperfunction, patients with TMJ pathology, lack of botulinum toxin therapy, muscle relaxants and / or tranquilizers.

At first, before treatment and during orthopedic structures setting, all the patients were tested using a device to determine the functional occlusion of T-Scan III Novus (Certificate of Registration №9887 / 2010, from 09.11.2010. Order of the Ministry of Health № 431, from 09.11.2010 UKTZED code: 9018 19 90 00) and apparatus for assessing the functional state of the masticatory muscles BioEMG III (Certificate of registration: №11216 / 2012, dated 10.02.2012. Order of the Ministry of Health: № 86, dated 10.02.2012 , UKTZED code: 9018 49 90 00. Manufacturer's name: BioResearch 42 Associated Inc., USA: Dental devices BioPac - register of medical equipment and medical devices). Determination of occlusion parameters was performed with the help of T-Scan III device, which defines and analyzes the force of tooth compression using ultra-thin sensors. T-Scan III technology is designed to perform dynamic determination of occlusion at all the stages of dental treatment and it is the only quantitative method of occlusion analysis used in practice. Diagnosis with the T-Scan III is very simple and painless. To determine the occlusal force, the patient must bite an individual plate – an ultra-thin sensor. At this time, all micromovements of the teeth are displayed on the computer monitor. The technology provides quantitative data on the strength and time of compression of the dentition. The data are obtained on two- and three-dimensional active diagrams, with the help of which it is possible to identify the problem points necessary for the further formation of the perfectly balanced occlusion and to do it quickly and accurately.

The methodology of research on the BioEMG III device is based on placing disposable electrodes on the previously degreased skin, these electrodes already have some electrically conductive gel in the projection of the motor points of the studied muscles. The obtained results are processed on a personal computer using special software (BioPACK BioRESEARCHAss Inc., USA), which allows to visualize and save the patient data immediately and quickly.

The use of the T-Scan III device in combination with the BioEMG III myograph makes it possible to determine the relationship between the electrical potential of the muscles and the occlusal relationships of the teeth.

With the help of these devices and the computer analysis the following functional indicators were studied: index of symmetry of the anterior bundles of temporal muscles, index of symmetry of the masticatory muscles, torque of the mandible, statically stabilizing occlusal index, relative efforts of balance disturbance for more than 10 % regarding the left and right, indicators of closing and opening time.

According to the results of electromyographic examination, the following indices were calculated, they were suggested by Ferrario V. F., and now they are the basis of modern interpretation of data obtained by electromyography [16]:

1. Index of symmetry of the anterior bundles of the temporal muscles (POC TA):

POC TA = (Average amplitude of the left temporal muscle / average amplitude of the right temporal muscle) X 100%

Symmetry index of the anterior bundles of the temporal muscles, % 83 \le (%) \le 100

Result: value and sign, indicates the predominance of the right or left temporal muscle during the study;

2. Symmetry index of the masticatory muscles (POC MM):

ROC MM = (Average amplitude of the left masticatory muscle proper \ average amplitude of the right masticatory muscle proper) $X\ 100\%$

Chewing muscle symmetry index, % 83 \le (%) \le 100 Result: value and sign, indicates the predominance of the right or left masticatory muscle during the study;

3. Torque of the lower jaw (TORS):

TORS = (Amplitude average of the left temporomandibular muscle + Amplitude average of the right proper masticatory muscle) \ (Amplitude average of the right temporal muscle + Amplitude average of the left proper masticatory muscle) X 100%

Lower jaw torque, % 90≤ (%) ≤100

Result: value, indicates possible displacement of the mandible horizontally when opening the mouth due to hyperactivity of the masticatory muscles;

4. Static stabilization occlusive index (BAR):

BAR = (Average amplitude of the left temporomandibular muscle + average amplitude of the right temporal muscle) / (average amplitude of the right masticatory muscle + average amplitude of the left masticatory muscle) X100%

Static stabilizing occlusal index, % 90 \le (%) \le 100

Result: value and sign, indicates the predominance of masseterial or temporal type of chewing (horizontal means masseterial, vertical – temporal).

According to the results of digital analysis of occlusiograms, the following indices were registered: relative efforts of balance disturbance for more than 10 % regarding the left and right, indicators of closing and opening time.

Patients were divided into two equal groups – the main group (patients were treated by using the suggested method of construction of occlusal surfaces of orthopedic structures) and control (dentition defects were replaced according to the traditional protocol). Orthopedic constructions for patients of the main group were performed on the basis of provisional constructions that met the needs of restoring proper articulation, muscle balance and muscle activity. Patients underwent construction of occlusal surfaces of orthopedic structures under monitoring of the dynamics of functional indicators of occlusion and the functional state of the masticatory muscles with subsequent correction of the dynamic component of these parameters. The comparison group included 15 people with intact dentitions. The study groups were randomized by sex, age, type of dentition defect.

It should be taken into consideration that in both the main and control groups, all the orthopedic structures were manufactured using the facial arch and individual

Table 1. Monitoring parameters of electromyographic examination in the process of orthopedic treatment (M \pm SE)

| Test groups | | Symmetry index of the anterior bundles of the temporal muscles, % | Symmetry index of the of massicatory muscles, % | Torque of the lower jaw, % | Statically stabilizing occlusive index % |
|----------------------------|------------------|--|---|----------------------------|--|
| control | Before treatment | 59 ± 5» | 70 ± 5» | 62 ±5» | 66 ±5» |
| n= 32 | After treatment | 75 ± 2* | 80 ± 2* | 76 ±2* | 70 ±2* |
| Main group n= 32 | Before treatment | 58 ± 5» | 71 ± 5» | 61 ±5» | 66 ±5» |
| | Diagnostic tests | 91 ± 2 | 90 ± 2 | 90 ± 2 | 90 ± 2 |
| | After treatment | 92 ± 2 | 90 ± 2 | 92 ± 2 | 91 ± 2 |
| Compari-son group n= 15 | | 90 ± 6 | 91 ± 7 | 94±3 | 93±3 |

Note: *- p>0,05 against the control group;

Table II. Parameters of digital occlusiography monitoring in the process of orthopedic treatment (M± SE)

| Group | os of research | Relative efforts of balance disturbance for more than 10 % regarding the left and right, | Time indicators of jaw closing, C | Time indicators of jaw opening, C |
|---------------------------|---|--|-----------------------------------|-----------------------------------|
| | Before treatment | 25 ± 0,6» | 0,6 ± 0,43» | 0,68 ±0,8» |
| Control n= 32 | After treatment | 23 ± 0,9 * | 0,512 ± 0,4 * | 0,589± 0,3 * |
| Main group n= 32 | Before treatment | 25 ± ,05» | 0,59 ± 0,5» | 0,61 ± 0,5» |
| | Diagnostic tests on provisional constructions | 5,8 ± 0,7 | 0,4 ± 0,24 | 0,3 ± 0,29 |
| | After treatment | 5,2 ± 0,5 | 0,343 ± 0,3 | $0,32 \pm 0,2$ |
| Comparison group n= 15 | | 2,1 ± 0,55 | 0,305 ± 0,15 | 0,313 ± 0,11 |

Note: *- p>0,05 against the control group;

articulators for maximum accuracy and anatomical and physiological compliance. Statistical processing of the obtained digital data was performed using the computer program Statistica 8.0 (STA862D175437Q). For each sample set of observations (n), the arithmetic mean (M), the standard error of the mean (SE), the standard deviation (SD), and the 95% confidence interval of the true mean were calculated using the Student's t-test. The level of significance of the mean values in the independent samples (p) was calculated according to the distribution function of the Student's t-test: at p <0.05 – the difference is significant; p> 0.05 – the difference is not significant.

RESULTS

The results of the electromyographic study (Table I) showed that in the comparison group (without dentition defects) the symmetry index of the anterior bundles of the temporal muscles was $90\% \pm 6$, the symmetry index of the masticatory muscles was $91\% \pm 7$, the torque of the mandible $-94\% \pm 3$, statically stabilizing occlusal index $-93\% \pm 3$. These values of the studied parameters indicate balanced and regulate activity of the masticatory muscle complex.

Analysis of digital occlusiograms in patients of the comparison group (Table II) showed that the value of relative efforts

of balance disturbance for more than 10 % regarding the left and right was determined at 2.1% \pm 0.55, the time of closure of the jaws was 0.305 s. \pm 0.15, and the opening time indicator is 0.313 s. \pm 0.15. The obtained data indicate a functionally correct occlusal correlation relation of the dentition.

Monitoring the results of the initial functional state of the masticatory muscles and the correlation of dentition in patients of the control group with dentition defects (Kennedy class III), not complicated by dental deformities showed significantly different results (p <0.05) from the comparison groups, which showed clear violations of occlusal relations. The parameters of electromyographic and occlusiographic examination, as can be seen from tables I and II, were as follows: the index of symmetry of the bundles of the anterior temporal muscles was, on average, 35% less than that without deformations of the dentition, the index of symmetry of the masticatory muscles was 24% less, the torque of the mandible was 34% less, the static index stabilizing occlusal was 29% less, relative efforts of balance disturbance for more than 10 % regarding the left and right, was 11.9 times more, the time of closing the jaws was 1.2 times more, the time of opening the jaws was 2.2 times more (p < 0.05). It should be taken into consideration that the initial values of these indicators in the main group (Fig. 1) were proportional to those in the

^{«-}p>0,05 against the comparison group

^{«-} p>0,05 against the comparison group



Fig. 1. 30 years-old patient M's myography (the main group, before treatment)



Fig. 2. 30 years-old patient M's myography (the main group, on the provisional constructions)

control (p> 0.05), which makes it possible to compare them further, and probably differ from such comparison groups (p<0.05). In the main group, a comparative evaluation of the values of diagnostic samples on makeshift structures was carried out according to the studied parameters in relation to the initial values (Table I,II; Fig. 2,3). Thus, the index of symmetry of the anterior bundles of the temporal muscles decreased by 36%, the index of symmetry of the masticatory muscles – by 21%, the torque of the mandible – by 32%, the index of statically stabilizing occlusal – by 26%, relative efforts of balance disturbance for more than 10 % regarding the left and right, increased 4 times, the rate of closing the jaws – 0.6 times, the time of opening – 2 times (p < 0,05). It is worth mentioning that these values of

functional parameters reached values commensurate with those in the comparison group (p > 0.05).

The results of evaluating the effectiveness of the proposed method on the dynamics of functional indicators of myoand occlusiography showed that their values in the main and control groups after orthopedic treatment had significant differences (p <0.05). Namely: the index of symmetry of the anterior bundles of the temporal muscles dropped by 18.5%, the index of symmetry of the masticatory muscles reduced by 22%, the torque of the mandible reduced by 32%, the index of stabilizing occlusal – by 27%, relative efforts of balance disturbance for more than 10 % regarding the left and right reduced by a factor of 4.4, the rate of opening time was 1.5 times higher than before, the rate of time of opening of the jaws – 1.8 times.




Fig. 3. 30 years-old patient M's digital occlusiography (the main group, on the provisional constructions,)




Fig. 4. 30 years-old patient M's digital occlusiography (the main group, after treatment)

The obtained results indicate the normalization of functional parameters when using the proposed method of constructing occlusal surfaces immediately after treatment (Fig. 4).

DISCUSSION

Until now, in orthopedic treatment, methods of restoring masticatory efficiency have taken into account only the qualitative indicators that characterize the condition of the dental-maxillary system. Quantitative monitoring of the dynamics of functional indicators of occlusion of T-Scan III Novus and the functional state of the masticatory muscles BioEMG III with subsequent correction of the dynamic component of these parameters allowed to achieve a positive clinical effect, which was manifested by the normalization of these indices.

This goal was achieved through the application of the proposed method of construction of occlusal surfaces of makeshift structures. The study of the condition of occlusal relations in patients with included defects of the dentition is reasonable and is preconditioned by the high prevalence and complexity of the diagnosis of functional disorders. The combined usage of myo- and occlusiography methods at different stages of orthopedic treatment is considered appropriate and promising.

The study of the condition of occlusal relations in patients with included defects of the dentition is reasonable and it is preconditioned by the high prevalence and complexity of the diagnosis of functional disorders. The combined usage of myo- and occlusiography methods at different stages of orthopedic treatment is considered appropriate and promising.

CONCLUSIONS

- 1. The combination of applied diagnostic methods of myo- and occlusiography makes it possible to assess the functional parameters of the masticatory apparatus in an objective way.
- 2. The use of the suggested method of orthopedic treatment with temporary structures leads to probable changes in the studied functional parameters in comparison with the control group (p < 0.05).
- 3. The obtained values of myo- and occlusiography indicators after the proposed treatment are commensurate with those in the comparison group (without dentition defects), which proves their normalization.

REFERENCES

- Nespriadko V.P., Moroz Yu.lu., Lyseiko N.V. Stan okliuziinykh spivvidnoshen u period adaptatsii do neznimnykh zubnykh proteziv [State of occlusion relations in the period of adaptation of patients to non-removable dentures]. Bukovynskyi medychnyi visnyk. 2017;2(82):42-46. (in Ukrainian).
- 2. Moroz lu.lu., Nespriadko V.P., Lusiuk S.V. Dynamyka yzmenenyia sostoianyia zhevatelnukh mushts y okkliuzyonnukh sootnoshenyi zubnukh riadov v rannye y otdalennue sroky protezyrovanyia patsyentov nes`emnumy zubnumy protezamy [Dynamics of changes in the state of the masticatory muscles and occlusal ratios of the dentition in the early and late periods of prosthetics of patients with fixed dentures]. Moderni vēda. 2018;1:112-120. (in Ukrainian).
- 3. Bida O. V. Patolohichni zminy okliuzii, obumovleni chastkovoiu vtratoiu zubiv, uskladnenoiu zuboshchelepnymy deformatsiiamy [Restoration of normal occlusal relationships in the complex treatment of patients with generalized periodontitis with included dentition defects]. Visnyk stomatolohii. 2016;4: 34-37. (in Ukrainian).
- Zhehulovych Z.Ye. Klinichna kharakterystyka dentalnoi okliuzii pislia vidnovlennia u konformatyvnomu pidkhodi [Clinical characteristics of dental occlusion after recovery in the conformational approach]. Novyny stomatolohii. 2015; 2(83): 18–23. (in Ukrainian).
- Fedorova O.V. Mozhlyvosti vykorystannia systemy komp'iuternoho analizu okliuziinykh kontaktiv pry ortopedychnomu likuvanni patsiientiv iz vtorynnymy zuboshchelepnymy deformatsiiamy (ohliad literatury) [Possibilities of the use of the system of computer analysis of occlusal contacts during orthopedic treatment of patients with secondary dento-facial deformations (review of literature)]. Ukrainskyi stomatolohichnyi almanakh. 2017;3: 49–52. (in Ukrainian).

- Kerstein R. B. Handbook of Research on Computerized Occlusal Analysis Technology Application in Dental Medicine. Global Book series. AMTCP. 2015.
- Bida O. V. Patolohichni zminy okliuzii, obumovleni chastkovoiu vtratoiu zubiv, uskladnenoiu zuboshchelepnymy deformatsiiamy [Pathological changes of occlusion caused by partial losses of teeth, complicated by teeth deformations]. Visnyk stomatolohii. 2016; 4: 34–37. (in Ukrainian).
- 8. Afrashtehfar K.I., Qadeer S. Computerized occlusal analysis as an alternative occlusal indicator. Cranio. 2016;34:52-57. doi: https://doi.org/10.1179/2151090314Y.0000000024
- 9. Klineberg I., Eckert S. Functional Occlusion in Restorative Dentistry and Prosthodontics. 1st ed. Mosby, 2016: 288.
- 10. Nelson S. Wheeler's Dental Anatomy, Physiology and Occlusion. 11th ed. Elsevier. 2019: 336.
- 11. Okeson JP. Management of Temporomandibular Disorders and Occlusion. 8th ed. Mosby. 2019: 51.
- Amin K., Vere J., Thanabalan N., Elmougy A. Occlusal concepts and considerations in fixed prosthodontics. Primary Dental Journal. 2019; 8 (3): 20-27.
- 13. Cerna M., Ferreira R., Zaror C., Navarro P. Validity and reliability of the T-Scan III for measuring force under laboratory conditions. The Journal of Oral Reabilitation. 2015; 42: 544-551.
- 14. Buduru S., Mesaros A., Talmaceanu D., Baru O. et al. Occlusion in the digital era: a report on three cases. Medicine and Pharmacy Reports. 2019; 92 (3): 78-84.
- 15. Afrashtehfar K. I., Qadeer S. Computerized occlusal analysis as an alternative occlusal indicator. Cranio. 2016; 34 (1): 52-57.
- Drohomyrets'ka M. S., Bilous M. K., Kushpela Y.I., Voytovych O. A. Craniopostural adaptation in orthodontic patients. Zbirnyk naukovykh prats' spivrobitnykiv NMAPO imeni P. L. Shupyka. 2015; 24 (1): 505–509.

ORCID and contributionship:

Svitlana P. Yarova:0000-0002-7830-4579 ^C Sergii O. Turchenenko:0000-0002-0396-357X ^{A, E,} Iryna M. Tkachenko: 0000-0001-8243-8644 ^{D, F} Yuriy Yu. Yarov:0000-0002-2867-9866 ^B

Conflict of interest:

The Authors declare no conflict of interest.

CORRESPONDING AUTHOR

Irvna M. Tkachenko

Ukrainian Medical Stomatological Academi 23 Shevchenko st., 36011 Poltava, Ukraine tel: +380505353181 e-mail: tkachenkoirmix@gmail.com

Received: 09.07.2020 **Accepted:** 20.11.2020

A-Work concept and design, B-Data collection and analysis, C-Responsibility for statistical analysis,

 ${f D}$ — Writing the article, ${f E}$ — Critical review, ${f F}$ — Final approval of the article